

Summary of NCD Alliance recommendations on proposed global targets for NCDs

	Proposed Target	WHO Best Buy	Highest WHO Criteria	NCD Alliance recommendations to Member States
1	Mortality from NCDs 25% relative reduction in overall mortality from cardiovascular disease, cancer, diabetes, or chronic respiratory disease		✓	Retain Endorse as overarching goal at World Health Assembly
2	Tobacco smoking 30% relative reduction in prevalence of current tobacco smoking	✓	✓	Retain
3	Dietary salt intake 30% relative reduction in mean adult (aged 18+) population intake of salt, with aim of achieving recommended level of less than 5 grams per day	✓	✓	Retain
4	Blood pressure /Hypertension 25% relative reduction in prevalence of raised blood pressure		✓	Retain
5	Physical activity 10% relative reduction in prevalence of insufficient physical activity in adults aged 18+ years	✓		Retain
6	Availability of essential medicines and technologies A minimum of 80% availability of affordable, quality-assured essential NCD medicines and technologies in public and private sectors			Promote the proposed indicators to this target
7	Prevention of heart attack and stroke 80% coverage of multidrug therapy (including glycaemic control) for people aged 30+ years with a 10 year risk of heart attack, stroke or diabetes ≥ 30%, or existing cardiovascular disease	✓		Reinstate
8	Alcohol 10% relative reduction in persons aged 15+ alcohol per capita consumption (APC)	✓	✓	Reinstate
9	Elimination of industrially produced trans-fats from the food supply	✓	✓	Reinstate
10	Obesity: No increase in obesity prevalence Childhood obesity: Sustained downward trend in prevalence of obesity in below-5s and school-aged children to ≤ 5% of the population by 2025			Reinstate ... or consider replacing with this target on childhood obesity

Summary of NCD Alliance recommendations on proposed global indicators for NCDs

Proposed Indicator	NCD Alliance recommendations
Unconditional probability of death between ages 30 and 70 years from cardiovascular diseases, cancer, diabetes, and chronic respiratory diseases.	Retain
Age-standardized prevalence of current tobacco smoking among persons aged 15+ years	Retain
Age-standardized prevalence of insufficiently active adults aged 18+ years (defined as less than 150 minutes of moderate-intensity activity per week, or equivalent) (%)	Retain